



Overview of Transport, Fast Particle and heating and current-drive Physics using Tritium in JET plasmas

D Stork,

K-D Zastrow, Yu Baranov, P Belo, L Bertalot, D Borba, J H Brzozowski, D Ciric, C D Challis,
S Conroy, M de Baar, P de Vries, P Dumortier, L Garzotti, N C Hawkes, T C Hender, E
Joffrin, T T C Jones, V Kiptily, J Mailloux, D C McDonald, M F F Nave, R Neu, M G
O'Mullane, J Ongena, R J Pearce, S Popovichev, S E Sharapov, M Stamp, J Stober, E Surrey,
M Valovič, I Voitsekhovitch, H Weisen, A D Whiteford, L Worth, V Yavorskij




































The JET "Trace Tritium" campaign

- The JET Trace Tritium Experiment (TTE) took place in September-October 2003.
- 20 dedicated experiments in 6 weeks
- Tritium present in plasmas in 'trace' quantities
 n_T/(n_T+n_D) ≤3%.
- Tritium introduced by short T₂ gas puffs (≤6 mg), or T⁰ beam injection (~100-300 ms 'blips')



Data derived from S/S 2.5:14 MeV neutrons

During TTE the **wall tritium fraction** kept at <0.5% by frequent pure **DD cleanup pulses**.

Ensured minimisation of background effects - enabled accurate quantification of tritium source







Physics with Trace Tritium

- Thermal particle transport
 - Diffusion (D_T) and convection (v_T) scaling and systematics including $\rho*,v*,\beta$ scaling studies (McDonald—EX/6-6);
 - Fuelling;
 - Effects of MHD phenomena on particle transport.
- Fast particle physics
 - Beam ion transport in Current Hole plasmas;
 - Fusion product confinement/ transport (Yavorskij –TH/P4-49)
- Neutron diagnostic development (Murari-- OV/P4-9)
- RF heating physics (Lamalle-- P5-165)

All TTE experiments were non-perturbative (D_T and V_T measured in plasmas where all relevant global and local background parameters were in steady-state).







Thermal Tritium Transport

• D_{τ} and v_{τ} can be measured separately. (unlike pure steady state deuterium $\Gamma = -D \frac{\partial n}{\partial r} + vn$ state deuterium

$$\frac{\partial n}{\partial t} = -\frac{1}{r} \frac{\partial}{\partial r} (r\Gamma) + \sigma(t)$$


$$\Gamma = -D \frac{\partial n}{\partial r} + vn$$

plasmas where only D_D/v_D is evaluated

- Accurate timing/ time profile of source $(T_2 \text{ gas} \sim 10 \text{ms}; \text{ thermalised NBI similar})$
- Evolution of tritium profiles derived from 14 MeV Neutron profiles

JET neutron profile monitor

- absolutely calibrated for 2.5 & 14 MeV neutrons.
- •19 channels at r/a<0.85
- •10 ms time resolution









Particle transport in many regimes

- Internal Transport Barrier (ITB) plasmas
- Hybrid scenarios $(q_0 \sim 1, +ve shear, no sawteeth)$
 - effect of plasma current and triangularity
- ELMy H-modes
 - scans of dimensionless parameters $\rho*, v*, \beta$;
 - variation of density;
 - effect of sawteeth;
 - [effect of heating profile(RF); impurity seeding].
- MHD effects --Neoclassical Tearing Modes (NTMs)
- Fuelling

In practically all regimes, particle transport found to be very much higher than neo-classical:

(except ITB region and high n_e near Greenwald density)







ITB particle transport


Pulse 61352 I_p=2.5 MA;


Reduction of D_T to neoclassical in the barrier region

 $B_{T} = 3.2 \text{ T}$

Otherwise diffusion>>neoclassical

ITB reduces v_T , but still stronger than neoclassical












ELMy H-mode particle transport

Greenwald density shot at $I_p=2.5$ MA

- ELMy H-mode results are averaged over ELMs and sawteeth.
- Diffusion is found to be strongly anomalous at low density:

as $n_e \sim n_{GREENWALD}$ then D_T approaches $D_{NEO,T}$ for r/a < 0.6

• Thermal diffusion varies less strongly with density than particle diffusion: thus $\chi_{\text{eff}}/D_{\text{T}}$ is density dependent.









ELMy H-mode particle transport:

Dimensionless parameter scans

- Each of $\rho*, v*$ (~naq/ T_e^2) and β were scanned in discharge pairs with T_2 gas puffs:
 - remaining two parameters held constant over the scan;
 - particle transport and energy transport results compared

McDonald :EX/6-6

- Inner (r/a<0.6) plasma shows Gyro-Bohm particle transport $D_T/B_0 \propto \rho^{*3.2}$
- Outer (r/a>0.6) plasma shows Bohm particle transport $D_T/B_0 \propto \rho^{*1.9}$
- β and $\nu*$ tritium transport fits contrast with Energy confinement for plasma r/a<0.6 particles: $D_{T,inner}/B_0 \propto \beta^{-0.34}$ and $\propto \nu^{*-0.51}$ for plasma r/a>0.6 particles: $D_{T,outer}/B_0 \propto \beta^{-0.55}$ and $\propto \nu^{*-0.4}$
- whereas energy confinement is largely independent of β and decreases weakly with $\nu*$








Thermal particle transport:

Systematics and ITER operational scenario comparison

Systematics between ITER scenarios are best compared outside the central region --no sawteeth or reversed shear etc.



- •Inside the sawtooth inversion radius, we expect differences between all three scenarios
- •Inside the ITB footpoint we expect differences in ITB discharges
- •We should see consistent behaviour for all three scenarios at r/a > 0.6








Thermal particle transport:

Systematics and ITER operational scenario comparison



- Strong correlation in all scenarios for D_T vs q_{95} .
- Try ${\rho_{pol}}^* = q \times \rho *$ as ordering parameter.
- ELMy H-modes show \sim Gyro-Bohm scaling in ${\rho_{pol}}^*$.
- Also see 1/β dependence when Hybrid scenarios are included
- Note also stronger convective v_T in ITBs.

Multiparameter fit
(NBI ELMy H-modes only)

 $D_{T}/B_{0} = c.\rho *2.4.q^{1.8}.\beta^{-1.08}.v*^{-0.13}$








Particle transport with NTMs

Otherwise identical discharges set up with and without Neoclassical Tearing Modes. Tritium gas puffs injected during the NTM period and evolution of the 14 MeV neutron profiles observed.










Particle transport with NTMs

Height (m)

Otherwise identical discharges set up with and without Neoclassical Tearing Modes. Tritium gas puffs injected during the NTM period and evolution of the 14 MeV neutron profiles observed.

NTM transport effects are confined to within the q=3/2 surface.










Particle transport with NTMs

NTM transport effects are confined to within the q=3/2 surface.

Shots with NTM show more rapid transit of Tritium to the plasma centre, and a quicker decay of tritium profile.

Effects can be explained by: higher D_T within q=3/2; or larger inward v_T at q=3/2 surface.



D Stork: TTE Overview - 20th Fusion Energy Conference - Vilamoura, Nov 2004





In the TTE Campaign, Fast particle physics results were obtained either:

- using NBI tritons directly
 - Beam ion transport in Current Holes;
 - [Transport of beam ions at low-q];
- using T⁰ NBI-derived fusion products
 - Fusion product confinement/ transport —
- [detecting effects of RF-accelerated protons pT fusion]







NBI transport in Current Holes

- Current Hole (CH) plasmas have near-zero Toroidal current density in a region with r/a < 0.3 0.5
- typically an ITB at the edge of CH region.
- Good thermal energy confinement inside the ITB, but CH expected to confine Fast Ions poorly due to low-central poloidal field.
- 105 keV Tritons injected into Monotonic Current profiles (MC) and CH plasmas, CH effects demonstrated by 14 MeV neutron profiles from interactions of fast tritons with background deuterium








NBI transport in Current Holes

14 MeV neutron profiles Data/simulations for on-axis NBI tritons



Outward profile shift seen in CH plasmas compared to MC plasmas Behaviour (due to outward movement of 'stagnation orbits' of the tritons) successfully modelled by 3-D Fokker Planck code and by TRANSP Monte Carlo







Fusion product transport/confinement (Yavorskij -TH/P4-49)


 γ -ray spectrometry has provided information on distribution function of fusion α - particles in TTE

 α -particle diagnosis was based on nuclear reaction ${}^{9}\text{Be}(\alpha, n\gamma)^{12}\text{C}$

following α -production in

$$T_{(F)} + D > n + \alpha$$

4.44-MeV γ -ray emission measures changes in the LOS density of the fast α -particles with E_{α} > 1.7 MeV post-NBI.










Fusion product transport/confinement (Yavorskij -TH/P4-49)

 γ -ray spectrometry has provided information on distribution function of fusion α - particles in TTE

 α -particle diagnosis was based on nuclear reaction ${}^{9}\text{Be}(\alpha, n\gamma)^{12}\text{C}$

following α -production in

$$T_{(F)} + D > n + \alpha$$



 γ -ray intensity after the T-blip is seen as $\sim I_{\gamma}(t) \propto \exp(-t/\tau_{\gamma})$







Fusion product transport/confinement (II)

For a wide range of discharges compare two quantities


- τ_γ measured 4.44-MeV
 γ-ray decay rate
- τ_T + τ_α calculated classical combined slowing down parameter

tritons from 105 keV to 40 keV plus

 α 's from 3.5+ MeV to 1.7 MeV

$$\tau_{\gamma}^{exp} \ge \tau_{\alpha} + \tau_{T}$$

$$\tau_{\gamma}^{exp} < \tau_{\alpha} + \tau_{T}$$









Fusion product transport/confinement (III)

$$\tau_{\gamma}^{exp} \ge \tau_{\alpha} + \tau_{T}$$

Good α -confinement: Critical plasma current $I_{CR} > 1.5-2MA$ to avoid significant orbit losses/drift


Discharges with monotonic *q*-profiles: $B_T = 2.2-3.2T$, $I_p = 2.0-3.0MA$

$$\tau_{\gamma}^{exp} < \tau_{\alpha} + \tau_{T}$$

Poor α -confinement: orbit losses and drift

Low plasma current : 1MA

'Current Hole' Discharges









Conclusions

- Turbulence dominates thermal particle transport for most regimes
 - Large inward ν_{τ} correlates with high D_{τ}
 - Neo-classical only for : high n_e ELMy H & in ITBs.
- Dimensionless parameters scans show:
 - Gyro-Bohm particle transport ($D_T \sim \rho^{*3}$) for Inner plasma;
 - Bohm particle transport ($D_T \sim \rho^{*2}$) for Outer plasma;
 - but when q scans are included behaviour is more like Gyro-Bohm in outer plasma ($D_T \sim \rho_{POI}^{*3}$, where $\rho_{POI}^{*} = q \times \rho^*$);
 - particle transport has an inverse β and ν * dependence.
- Redistribution of fast deuterium and thermal tritium by sawteeth and NTMs is observed
- Current Hole plasmas show effects on Fast Ion transport and confinement, which can be modelled qualitatively by 3-D Fokker Planck code or TRANSP.







Reserve slides

Hybrid scenarios








Hybrid scenario results

- T₂ gas puff and T⁰ NBI were injected into 'Hybrid scenario' plasmas [Joffrin -EX/4-2]
 - triangularity scan (δ =0.2 -0.46) at fixed confinement ($H\beta_N/q_{95}^2 \sim 0.32$, $H\beta_N \sim 5-5.5$);
 - I_{P} (1.4 MA -> 2.0 MA) scan at constant q, and low δ
- A global particle confinement time (τ_p) can be established approximately from the decay of the 14 MeV neutrons after thermalisation of fast NBI-ions



- The effect of wall recycling on this value of τ_P is negligible
- Global τ_{P} increases by ~50% over the δ scan and scales ~ with I_{P}





Hybrid scenario results (II)

- Global τ_P increases by ~50% over the δ scan and scales ~ with I_P .
- The fits to the 14 MeV profiles maintain this result quantitatively.
 - For triangularity scan, difference appears as reduction of D_T across the whole plasma, as δ ↑.
 - For current (not shown) there is a less significant reduction in D_T for r/a > 0.5 as I_P increases

