Review of D-T Experiments Relevant to Burning Plasma Issues

R. J. Hawryluk

Frontiers in Plasma Confinement and Related Engineering/Plasma Science

Toki, Japan

December 11, 2001

PPPL
PRINCETON PLASMA PHYSICS LABORATORY

Excellent Technical Progress Has Enabled TFTR and JET to Begin Studying Burning Plasmas



JET and TFTR were Designed to Perform Deuterium-Tritium Experiments

Joint European Torus (JET)



Inside the Joint European Torus (JET) vacuum vessel, JET Joint Undertaking, UK

Tokamak Fusion Test Reactor (TFTR)



	JET	TFTR
I_p(MA)	4.0	2.7
B_T(T)	3.6	5.6
P_{DT}(MW)	16.1	10.6
$P_{DT}/P_{in}=Q_{DT}$	0.64	0.27
Fusion Energy (GJ)	0.68	1.70
Tritium Proc. (g)	99.3	99.

How are D-T Experiments Different from D?

- **Isotope Effects**
  - **Transport**
  - **Wave-particle Interactions**
- **Alpha-particle Physics**
- **Technology**

PPPL
PRINCETON PLASMA PHYSICS LABORATORY

Isotope Effect Studied in a Wide Variety of Operating Regimes



- **Challenge to theoretical interpretation and to gyro-Bohm scaling, $\langle A \rangle^{-0.2}$**
- **Recent ITER scaling $\langle A \rangle^{0.19}$**

JET D-T Elmy H-mode Experiments Consistent with ITER Scaling





- **What is the role of the pedestal and the core?**

- $\tau_{th} \propto \langle A \rangle^{0.16 \pm 0.06}$

Stored Energy Associated with Pedestal Increases with <A>



- $W_{ped} \propto \langle A \rangle^{0.96}$
- Power loss by ELMs decreases

In JET H-mode Discharges Thermal Conductivity Degrades in the Core





- **Confinement scaling in the core consistent with gyro-Bohm.**
  - $\tau_{thcore} \propto \langle A \rangle^{-0.16 \pm 0.1}$

    $\chi_I \propto \langle A \rangle^{0.73 \pm 0.4}$

Power Threshold Going from L to H-mode Shows Favorable Isotope Scaling





- $P_{loss} \propto \langle A \rangle^{-1}$
- ELM frequency decreases with <A>

PPPL PRINCETON PLASMA PHYSICS LABORATORY

Improvement of χ_I in Core of TFTR Supershots in DT





$\tau_E^{thermal} \propto \langle A \rangle^{0.89\pm0.1}$

$\chi_i^{tot} \propto \langle A \rangle^{-1.8\pm0.2}$

- $n_i(0)\tau_E T_i(0)$ increased by ~55% from D to DT
  - Some cases up to 80% increase

Scott

ITG Model with Radial Electric Field Reproduces the Ion Temperature

TFTR



- **Maximum linear growth rate decreases with ion mass.**
- **Er shearing rate increases with Ti.**
- **Radial Electric Field Shear Reproduces Strong Isotope Effect.**
- **==> Isotope Effect depends on Operating Regime**

Ernst

PPPL PRINCETON PLASMA PHYSICS LABORATORY

Favorable Isotope Scaling Not Observed in Reverse Shear Experiments



- **Power threshold for internal barrier formation increased with <A>.**

Summary of Isotope Effects on Confinement

- H-mode isotope scaling studies on JET, together with the worldwide physics database, provide a good technical basis for baseline operation of burning plasma experiments.
  - Isotope scaling for τ_E and power threshold.

- Understanding of isotope scaling is incomplete.
  - Variation in different operating regimes.
  - What is the role of radial electric field shear?
  - What are the implications for advanced operating modes?

PPPL
PRINCETON PLASMA PHYSICS LABORATORY

ICRF Successfully Heated DT Supershot Plasmas in TFTR



ΔT_i due to 2nd harmonic tritium heating



ΔT_e due to direct electron and ^{3}He minority ion heating

- **Power deposition calculations in good agreement with experiment.**

JET Demonstrated Successful Deuterium Minority Heating of Tritium Plasmas



Eriksson, Start

- **Record Q_{DT} for steady-state operation.**
  - deuterium energy optimized
  - Not optimal for high Q_{DT}
- **Strong ion heating observed with ^{3}He**
  - Absorption weaker with $2\Omega_T$
  - Recommended scenario for ITER
- **Studied tritium minority heating.**

- **Physics of ICRF heating well established.**
  - Technology challenges remain.

PPPL PRINCETON PLASMA PHYSICS LABORATORY

Alpha-particle Physics Studies

- **MHD Quiescent**
  - **Alpha-particle heating**
- **MHD Affects Alpha Confinement**
- **Alpha Particle Induced MHD Activity**
- ***Diagnostic Development Critical for these Studies***

PPPL
PRINCETON PLASMA PHYSICS LABORATORY

Alpha-Particle Parameters in TFTR/JET Sufficient to Begin Study of Alpha-Particle Physics

	TFTR	JET	ITER
P_{fusion}(MW)	10.6	16.1	400
$p_\alpha(0)$ (MW/m^3)	0.28	0.08	0.43
$\beta_\alpha(0)$%	0.30	0.4	0.8
$-R\bullet grad(\beta_\alpha)$%	2.0	2.3	4.0
$V_\alpha(0)/V_{Alfvén}(0)$	1.72	2.52	1.76
P_α/P_{heat}	0.03	0.09	0.66

Budny

PPPL PRINCETON PLASMA PHYSICS LABORATORY

D-T Neutron Emission is Consistent with Calculations Based on Plasma Parameters



Budny, Keilhacker



- **Alpha birth rate and profile are adequately modeled.**

Escaping Alpha Flux at 90° Detector is Consistent with Classical First Orbit Losses



- **At 2.5 MA in TFTR , first orbit loss ≈ 3% globally**

Confined Alphas in the Plasma Core Show Classical Slowing Down Spectrum

TFTR



Double Charge Exchange Technique

$He^{++} + Li^{+} \Rightarrow He^{0} + Li^{3+}$



- **TRANSP calculation includes:**
  - **orbit trajectories**
  - **classical slowing down**
  - **time dependence of alpha production**

Fisher, Petrov, Medley

PPPL PRINCETON PLASMA PHYSICS LABORATORY

Stochastic Ripple Diffusion Affects Confinement of Deeply Trapped Particles



- **FPPT includes modeling of stochastic ripple diffusion.**
- **Heat deposition due to ripple loss of fast ions imaged on JT60U.**

Alpha Particles Are Well Confined





Charge exchange between fast beam neutrals and nonthermal alphas

- **$0 \leq D_\alpha \leq 0.03$ m²/s**

McKee

Rapid Ash Transport from the Core to the Edge in Supershots

TFTR



- Consistent with $\tau_p^{*He}/\tau_E = 8$, acceptable for a reactor

$D_{He} / \chi_D \sim 1$

Synakowski

- **Confinement of ash consistent with impurity transport and edge conditions.**

PPPL PRINCETON PLASMA PHYSICS LABORATORY

Sawteeth Cause a Large Radial Redistribution of Alpha Particles

TFTR



Stratton

- **Alpha heating profile in a burning plasma will depend upon sawtooth activity, although only transiently**

MHD Activity can Cause Enhanced Loss of Alpha Particles

TFTR



Zweben, Darrow

- Strong toroidal anisotropy in loss apparent as NTM mode is rotating
- Enhanced loss observed due to:
  - disruptions
  - kinetic ballooning modes
- Concern for plasma-facing components.

PPPL PRINCETON PLASMA PHYSICS LABORATORY

Observation of TAE Instability Driven by Fusion Alphas





Radial mode structure via microwave reflectometry.

Nazikian

PPPL PRINCETON PLASMA PHYSICS LABORATORY

TAE Modes Redistribute Deeply Trapped Alpha-Particles



Observed
q(0) ~ 2

- Further work required to benchmark models.
- NNBI experiments on JT60U have observed strong radial redistribution.

Sawtooth Frequency Decreases with Isotopic Mass





- **Change in sawtooth period attributed to increase in slowing down time and beam ion perpendicular energy density.**
- **Calculations imply that alpha particles had a stabilizing effect in highest performance D-T discharges.**

PPPL PRINCETON PLASMA PHYSICS LABORATORY

Initial Evidence of Alpha-Particle Heating



TFTR

- **Alpha heating ~15% of power through electron channel**
- **Plasmas matched for dominant Te scaling in D only plasmas.**
  - **$T_e(0) \propto \tau_E^{0.5}$**
  - **account for the isotope effect on confinement**
- **No sawteeth or adverse MHD.**

Taylor, Strachan

Alpha Heating Observed in JET



- **P_{alpha}/P_{heat} ~12%**
  - 30-40% through the electron channel
- **Larger range in T concentration**
  - analyze the effects associated with isotopes
- **Sawteeth variability**
  - limited number of pulses

- **Alpha heating observed on both TFTR and JET.**
- **Comprehensive study of alpha heating requires higher values of P_{alpha}/P_{heat} .**

Can Alpha Particles Be Harnessed to Further Improve Reactor Potential?

- **Theoretical work on alpha channeling, frequency sweeping of waves, and stochastic ion heating raise the prospects of:**
  - **Increased ion heating**
  - **Alpha ash control**
  - **Modify alpha heating profile**
  - **Reduce alpha pressure to decrease instability drive**
  - **Current drive**

- **Need for experimental study**
  - **Fundamental wave-particle physics studies begun on TFTR.**
  - **D. Gates et al. predicted the role of stochastic ion heating in NSTX discharges when $V_{NBI} >> V_{Alfven}$**
  - **P. Thomas at EPS suggested possibility of increased ion heating due to alpha particles.**
    - ***Limited data***

PPPL
PRINCETON PLASMA PHYSICS LABORATORY

Summary Alpha-Particle Physics

- Alpha-particle confinement, transport, and slowing down well understood in MHD quiescent discharges.
  - Not as extensively studied in reversed shear experiments.
- Alpha-particle confinement affected by MHD.
- Alpha particle driven instabilities studied
  - Nonlinear consequences remain to be studied.
- Alpha-particle heating observed.
  - Higher power alpha particle heating experiments are required.

PPPL
PRINCETON PLASMA PHYSICS LABORATORY

Tritium Processing Safely Performed

- On-site tritium processing successful.
- Tritium retention in graphite is a serious concern.
  - TFTR tiles 16% retention
  - JET 12% retention
  - One year after extensive removal efforts
- Need for in-situ removal
  - Alternative materials



PPPL-JAERI Collaboration

Gentile,Skinner (PPPL)

Shu (JAERI)

PPPL PRINCETON PLASMA PHYSICS LABORATORY

In-vessel Remote Handling Demonstrated on JET



Telescopic Articulated Manipulator

JET

- **Critical technology for future burning plasma experiments**
  - **Fluence ~ 10^4 greater**

PPPL PRINCETON PLASMA PHYSICS LABORATORY

Successful Demonstration of Decommissioning of a Fusion Facility

TFTR



- **Machine activation in future burning plasma experiments will require remote disassembly.**

PPPL
PRINCETON PLASMA PHYSICS LABORATORY

Summary

- **Results from TFTR and JET:**
  - **Provided solid design basis for a burning plasma experiment.**
  - **Identified opportunities to learn new science and technology.**
- **Critical aspects of the technology for a burning plasma experiment were utilized**
  - **Burning plasma experiment will be far more demanding due to the higher fluence, tritium retention and processing requirements.**
- ***Full potential and consequences of alpha heating have not been explored!***

PPPL
PRINCETON PLASMA PHYSICS LABORATORY