


***Fusion and Plasma Physics are at the Core of Nature's Most Powerful Self-Driven Systems***


---

***Eagle Nebula***



***HST***




***Cassiopeia A***



***CHANDRA***

# Can we Solve the Mystery of Producing a Stationary Self-Sustained Fusion Fire??

**Galactic Jet - M87**



---

**Confining a Fusion Fire**  
**A Grand Challenge for Science and Technology**

**Dale Meade**

**Princeton University**

**Presented at**

**Idaho Section of the ANS**

**INEEL, Idaho Falls, ID**

<http://fire.pppl.gov>

**May 23, 2002**


<http://www.cpepweb.org>

---

# Fusion is an Outstanding Physics Challenge and is Connected to Other Outstanding Challenges

---


December 1999



## Ten Outstanding Physics Challenges

- Quantum gravity presents the ultimate challenge to theorists
- Explaining high- $T_c$  superconductors
- Unstable nuclei reveal the need for a complete theory of the nucleus
- Realizing the potential of fusion energy
- Climate prediction is heavy weather
- Turbulence nears a final answer
- Glass physics: still not transparent
- Solar magnetic field poses problems
- Complexity, catastrophe and physics
- Consciousness: the physicists view




# Fusion Does Work at Large Size




**Why is it so difficult in the lab?**

## Relevant Reactions for Fusion in the Laboratory

---




# Fusion Cross Sections and Reaction Rates



**For Example:**

$$\begin{aligned}
 P_{DT} &= n_D n_T \langle \sigma v \rangle (U_\alpha + U_n) \\
 &= 5.6 \times 10^{-7} \langle \sigma v \rangle n_D n_T \text{ watts m}^{-3}, \text{ note: } \langle \sigma v \rangle \sim T^2 \text{ @ } 10 \text{ keV}
 \end{aligned}$$

# The Grand Challenge, Science and Technology for Fusion



## Key Plasma Performance Metrics


- **Fusion Gain ( $Q_p$ )**
- Fusion Energy Density
- Duty Cycle/Repetition Rate

## Key Engineering Metrics

- **First Wall Lifetime**
- Availability/Reliability
- Environment and Safety
- System Costs


# There are Three Principal Fusion Concepts

---




## Spherical Inertial

gravitational  
transient compression  
drive (laser-D/I, beam)  
radial profile  
time profile  
electrostatic



## Toroidal Magnetic

surface of helical B lines  
twist of helix  
twist profile  
plasma profile  
toroidal symmetry



## Reactivity Enhancement

muon catalysis  
polarized nuclei  
others?

# Plasma Requirements for a Burning Plasma

---

## Power Balance

$$P_{\text{aux-heat}} + n^2 \langle \sigma v \rangle U_{\alpha} V_p / 4 - C_B T^{1/2} n_e^2 V_p = 3nkTV_p / \tau_E + d(3nkTV_p) / dt$$

where:  $n_D = n_T = n_e / 2 = n / 2$ ,  $n^2 \langle \sigma v \rangle U_{\alpha} V_p / 4 = P_{\alpha}$  is the alpha heating power,  $C_B T^{1/2} n_e^2 V_p$  is the radiation loss,  $W_p = 3nkTV_p$  and  $\tau_E = W_p / (P_{\text{aux-heat}} - dW_p / dt)$  is the energy confinement time.

## In Steady-state:

$$n\tau_E = \frac{3kT}{\langle \sigma v \rangle U_{\alpha} (Q+5)/4Q - C_B T^{1/2}}$$

where  $Q = P_{\text{fusion}} / P_{\text{aux-heat}}$

$Q = 1$  is Plasma Breakeven,  $Q = \infty$  is Plasma Ignition



# Comparison of Typical Plasma Parameters for Inertial and Magnetic Fusion

---

	<u>Inertial</u>	<u>Magnetic</u>
$T_i$ (keV)	10	10
$n$ ( $m^{-3}$ )	$6 \times 10^{30}$	$3 \times 10^{20}$
$\tau_E$ (sec)	$10^{-10}$	2
radius (m)	$10^{-4}$	1

# Why is Confinement a Challenge for Magnetic Fusion?

---

A D-T reactor at a fuel density of  $10^{20} \text{ m}^{-3}$  requires

$$\tau_E \sim 1 \text{ second}, \quad T_i \sim 10 \text{ keV}$$

$$T_e \sim 10 \text{ keV} \quad v_{te} \sim 6 \times 10^7 \text{ m/s}$$

Assume a container with


radius  $\sim 1 \text{ m}$  (typical radius for a magnetic bottle)

Then the number of bounces

$$N \sim v_{te} / r \sim 6 \times 10^7$$

$\sim 30$  coulomb collisions under typical conditions

# Toroidal Magnetic Chamber (Tokamak)



## Axisymmetric Magnetic Configuration

- axial current,  $I_T$ , produces toroidal magnetic field,  $B_T$
- toroidal current,  $I_p$ , produces vector potential,  $A_\phi$  and poloidal field,  $B_p$

## Axisymmetry ensures that:


- magnetic field lines lie in nested magnetic surfaces given by  $\Psi = 2\pi R A_\phi$
- charged particles are confined to within  $\delta$  of magnetic surface due to conservation of canonical angular momentum

$$2mH = p_R^2 + p_Z^2 + \frac{(p_\phi - eRA_\phi)^2}{R^2} + e\Phi(R,Z)$$

$$\delta \sim mv/eB_p$$

# Toroidal Magnetic Confinement

---




Nested Magnetic Surfaces ( $\Psi = \text{constant}$ )

Plasma pressure =  $f(\Psi)$


$q = q(\Psi)$

Toroidal Asymmetry can cause plasma loss

- small magnetic field perturbations can have large effect at resonant surfaces
- particle collisions (would allow present tokamaks to be near ignition)
- **plasma instabilities (main limit in present fusion devices)**



**Tokamak Fusion Test Reactor**




**Tokamak Fusion Test Reactor**

# Comprehensive Diagnostic Systems have been Developed to Investigate Fusion Plasmas

## Spatially and Time Resolved

— Typically ~50 diagnostic measurements

- Equilibrium magnetics
- Core profile diagnostics  
 $n_e, n_i, T_i, T_e, Z_{\text{eff}}, Z_i, v$
- Internal magnetic field profile,  $B_\theta, q$
- Core and edge turbulence  $\tilde{n}_e, \tilde{T}_e$
- Edge and divertor  $T_e, n_e, Z_e, \text{radiation, neutral pressure}$



# Plasma Instabilities Limit Fusion Plasma Confinement

---

## Small-Scale Electrostatic Turbulence (fluctuating electric field, $E$ )

$\mathbf{v} = \mathbf{E} \times \mathbf{B} / B^2$ , ions and electrons both drift across the magnetic field preserving charge neutrality

$> \lambda_i$  instability wavelength  $\sim$  ion gyro-radius

$\sim V$  correlation random walk step size

## Small-Scale Magnetic Turbulence (fluctuating magnetic field, $B$ )

$\mathbf{v} = v_{\text{thermal}} \mathbf{B} / B$ , mainly loss of electron energy

## Large-Scale Large-Amplitude Magnetic Instability

plasma pressure sufficient to distort even tear the magnetic field, similar to solar flares. Can cause total loss of plasma in a tokamak.

# FUSION POWER IS DETERMINED BY MACROSCOPIC STABILITY

---


- **Plasma stability is largely determined by**

$$\beta \equiv \frac{2nT}{B^2 / 2\mu_0}$$

- **Fusion power**


$$P_{fus} = E_{fus} n_d n_t \langle \sigma_{fus} v \rangle \sim n^2 T^2 \sim \beta^2 B^4$$

- ***Denser, hotter plasma makes more fusion.***



# Simulation of a Plasma Disruption Driven by High Plasma Pressure

---




Nonlinear 3-D Fluid Computation

# Plasma Science Areas in Magnetic Fusion

---


- **Macroscopic Stability**
- **Wave-particle Interactions**
- **Transport and Microturbulence**
- **Plasma-wall Interactions**
- **Self-heated Plasmas**



# Wind Tunnel Experiments on Plasma Confinement


Dimensionless Parameters
$\omega_c \tau$
$\rho^* = \rho/a$
$v^* = v_c/v_b$
$\beta$

Similarity Parameter
$B R^{5/4}$



Kadomtsev, 1975

# Measured Transport is Much Larger than Neoclassical Transport




- Wrong profile, scaling with B and collisionality
- Better than no magnetic field by  $10^6$
- Additional processes: turbulence



# Localized Turbulence Measured via BES

- Beam Emission Spectroscopy: measure local density turbulence from fluctuations in light emitted from injected neutral  $H^0$  beam:


## BES Viewing Geometry



## Poloidal Cross Section



# Reconstruction of Spectral Data Showing Turbulent Eddies in TFTR



- $\delta n/n \sim 0.1 \%$ ,  $\delta T_i/T_i \sim 3-4 \delta n/n$ ,  $\lambda \gg \rho_i$ ,  $\lambda_{\text{radial}} \gg \lambda_{\text{poloidal}}$
- **Consistent with simulations of ion temperature gradient (ITG) instabilities**

# Understanding Turbulent Plasma Transport

---

## An **important problem**:


- ⌞ -- Size of plasma ignition experiment determined by fusion self-heating versus turbulent transport losses
- ⌞ -- Dynamics also of interest to other fields (e.g., astrophysical accretion disks)

⌞ A **scientific *Grand Challenge* problem**

⌞ A **true terascale computational problem for MPP's**

# Full Torus Simulations of Turbulent Transport Scaling

---





- Large-scale full torus gyrokinetic particle simulations for device-size scans
- Global *field-aligned mesh* (GTC code) saves factor  $\sim 100$  in computation
- Efficient utilization of new 5 TF IBM SP @ NERSC (just available 8/01) -- *fastest non-classified supercomputer in world*
- Most recent simulations used *1 billion particles (GC)*, *125 M spatial grid points*, and *7000 time steps* --- leading to important (previously inaccessible) new results

# Turbulent Fluctuations Suppressed When $E \times B$ Shearing Rate Exceeds Maximum Linear Growth Rate of Instabilities

TFTR


## Gyrokinetic Simulations

- Turbulent eddies disrupted by strongly sheared plasma flow



## Experiment

- Bursts of fluctuations are suppressed when  $E \times B$  shearing rate exceeds growth rate of most unstable mode




E. Mazzucato *et al*, PRL 77 (1996) 3145




# High Energy Physics Accelerators Enable Discovery

**HEP facilities plotted by discovery reach in mass vs. year**

**Also shown are some important discoveries and the expected range for the Higgs**




# Magnetic Fusion Pathways to Self-Heating



# Burning Plasma Physics - The Next Frontier


---

Three Options  
(same scale)




## FIRE

US Based  
Diversified International Portfolio



## ITER-FEAT

JA, EU or CA Based  
International Partnership




## IGNITOR

Italian Based  
International Collaboration

# Fusion Ignition Research Experiment

## (FIRE)

<http://fire.pppl.gov>




### Design Features

- $R = 2.14 \text{ m}$ ,  $a = 0.595 \text{ m}$
- $B = 10 \text{ T}$
- $W_{\text{mag}} = 5.2 \text{ GJ}$
- $I_p = 7.7 \text{ MA}$
- $P_{\text{aux}} \leq 20 \text{ MW}$
- $Q \approx 10$ ,  $P_{\text{fusion}} \sim 150 \text{ MW}$
- Burn Time  $\approx 20 \text{ s}$
- Tokamak Cost  $\approx \$375\text{M}$  (FY99)
- Total Project Cost  $\approx \$1.2\text{B}$  at Green Field site.

### Mission:

Attain, explore, understand and optimize fusion-dominated plasmas.

# Simulation of Burning Plasma in FIRE



- ITER98(y, 2) with  $H(y, 2) = 1.1$ ,  $n(0)/\langle n \rangle = 1.2$ , and  $n/n_{GW} = 0.67$
- Burn Time  $\approx 20 \text{ s} \approx 21\tau_E \approx 4\tau_{He} \approx 2\tau_{CR}$

$$Q = P_{\text{fusion}} / (P_{\text{aux}} + P_{\text{oh}})$$


## Concluding Remarks

---


- The capability now exists to produce and control fusion plasmas for detailed investigation in the laboratory. However, fusion reactors based on the present state of knowledge are large and innovations are needed for an attractive reactor concept.
- Recent developments in plasma diagnostics and computer simulation of three-dimensional non-linear phenomena now allow detailed comparison of theory and experiment.
- New insight into the physical processes causing plasma transport could lead to an advanced toroidal configuration that would have a significant impact on the attractiveness of magnetic fusion.
- The FIRE compact high field tokamak could address many of the generic fusion science issues including: self-heated plasma physics, many of the long pulse advanced tokamak issues and could begin the study of self-heated self-organized plasmas in a \$1B class experimental facility.

<http://fire.pppl.gov>

**Laboratories to Explore, Explain  
and Expand the Frontiers of Science**




**CHANDRA**



**VLBA**




**NIF**




**FIRE**



**HST (NGST)**



**SNS**



**CHANDRA**