

Gyrokinetic Studies of Turbulence Spreading


T.S. Hahm, Z. Lin,^a P.H. Diamond,^b G. Rewoldt,
W.X. Wang, S. Ethier, O. Gurcan,^b W. Lee, and W.M. Tang

Princeton University, Plasma Physics Laboratory, USA

^aUniv. of California, Irvine, USA

^bUniv. of California, San Diego, USA

Nov. 3, 2004, Vilamoura, Portugal




Outline and Conclusions

- Turbulence spreading into linearly stable zone is studied using global gyrokinetic particle simulations and theory.
- Motivation from experiments to study spreading of Edge Turbulence into Core.
- Results
 - Fluctuation amplitude in the linearly stable zone can be significant due to turbulence spreading.
 - Sometimes **Spreading of Edge Turbulence** into Core can exceed local turbulence in connection region.
 - It is likely to affect “the edge boundary conditions” used in core modeling, and predictions of pedestal extent.

Determination of Fluctuation Amplitude

$$\gamma = \gamma_{lin} - k_{\perp}^2 D_{turb} \rightarrow 0$$

- Nonlinear coupling induced dissipation leads to saturation (B. Kadomtsev '65)




- “Local Balance in Space” for a mode \mathbf{k}
- “Conceptual Foundation of Most Transport Models”
- **Missing:**
 - Meso-scale Phenomena: Barrier Dynamics, Avalanches,...
 - Anomalous transport in the region $\gamma_{lin} < 0$
 - **Turbulence Spreading into Less Unstable Zone**

Excitation of Linearly Damped Modes

- Nonlinear Saturation from Balance between:

γ_{lin} vs. **Spectral Transfer** from Nonlinear Mode Coupling



→ **Non-zero Amplitude for Linearly Damped Modes**

Sagdeev and Galeev, *Nonlinear Plasma Theory* (1969)

Gang-Diamond-Rosenbluth, *Phys. Fluids B* **3**, 68 (1991)


Hahm-Tang, *Phys. Fluids B* **3**, 989 (1991)

Horton, *Rev. Mod. Phys.* (2000) *for more references*

⇒ Lin *et al.*, IAEA/TH/8-4 (2004), this Friday

Nonlinear Coupling Leads To Radial Diffusion

- Nonlinear interactions of modes must spread fluctuation energy in radius due to:
 - i) $ik_x \rightarrow \frac{\partial}{\partial x}$
 - ii) poloidal harmonics at $q(r) = m/n$
 - iii) with different radial extents
 - iv) Numerical Studies with both Linear Toroidal Coupling and Nonlinear Coupling



[Garbet-Laurent-Samain-Chinardet, NF 1994]

- $\mathbf{E} \times \mathbf{B}$ nonlinearity \rightarrow “local turbulent damping” and “radial diffusion”:

$$(\mathbf{k} \times \mathbf{k}' \cdot \mathbf{b})^2 R_{k,k'} I_k I_{k'} \rightarrow -\frac{\partial}{\partial x} D_r(I) \frac{\partial}{\partial x} I + k_\theta^2 D_\theta(I) I.$$



[eg., Kim-Diamond-Malkov-Hahm *et al.*, NF 2003]

Simple Model of Turbulence Spreading

[Hahm, Diamond, Lin, Itoh, Itoh, PPCF 46, A323 '04]

$$\frac{\partial}{\partial t} I = \gamma(x)I - \alpha I^2 + \chi_0 \frac{\partial}{\partial x} \left(I \frac{\partial}{\partial x} I \right)$$



- $\gamma(x)$ is “local” growth rate, α : a local nonlinear coupling
- $\chi_0 I = \chi_i$ is a turbulent diffusivity
- I : turbulence intensity, $\Sigma_{\mathbf{k}} \text{ Modes} \sim \Sigma \text{ Eddys}$



$$\frac{\partial}{\partial t} \int_{x-\Delta}^{x+\Delta} dx' I(x', t) \sim \chi_0 I \frac{\partial}{\partial x} I \Big|_{x-\Delta}^{x+\Delta} + \dots$$


- **Profile of Fluctuation Intensity** crucial to its Spatio-temporal Evolution

Turbulence Spreading after Local Saturation




From Gyrokinetic (GTC) simulations, turbulence spreads radially ($\sim 25\rho_i$) into the linearly stable zone, causing deviation from GyroBohm scaling.

[Lin *et al.*, Phys. Rev. Lett. (2002)]




Propagation and Saturation of Fluctuation Front

- The **nonlinear** diffusion, in the absence of dissipation, will make the front propagate beyond x_0 indefinitely.



Front propagation stops when radial flux due to propagation is balanced by dissipation:



$$T_{prop} \simeq \Delta / U_x \iff T_{damp} \sim (|\gamma'| \Delta)^{-1}$$



$$\Delta^2 \simeq \frac{12\chi_0 I_0}{|\gamma'| x_0}, \text{ using the values from simulation } \rightarrow \Delta \simeq 18\rho_i$$

From GK simulation for a profile considered: $\Delta \simeq 25\rho_i$

Connection Region between Edge and Core



- Profile of Turbulence Intensity crucial in turbulence spreading: $\Gamma_I = -\chi(I) \frac{\partial}{\partial x} I$
- Core confinement improvement after L-H transition:
JET, ASDEX, DIII-D, C-mod,...
- **Connection Region:**
Local Turbulence + **Incoming Edge Turbulence**

Turbulence Spreading from Edge to Stable Core

- Nonlinear GTC Simulations of Ion Temperature Gradient Turbulence:

$\frac{R}{L_T} = 5.3$ at core
(within Dimits shift regime)


$\frac{R}{L_T} = 10.6$ at edge:

- Initial Growth at Edge
→ Penetration into stable Core
(Lin-Hahm-Diamond,
PRL '02, PPCF, PoP '04)

- Saturation Level at Core:


$$\frac{e\delta\phi}{T_e} \sim 3.6 \frac{\rho_i}{a}$$

$$\rightarrow \nabla \cdot \Gamma_I \gg \gamma_{local} I$$



Spreading in Unstable Zone

[Gurcan, Diamond, Hahm, and Lin, *Submitted to Phys. Plasmas* '04]



$$\frac{\partial}{\partial t}I = \gamma(x)I - \alpha I^2 + \chi_0 \frac{\partial}{\partial x} \left(I \frac{\partial}{\partial x} I \right)$$

- When $\gamma(x)$, α , χ_0 are constant in radius, the Fisher-Kolmogorov equation with nonlinear diffusion exhibits **“propagating front”** solutions.
- The spreading can beat local growth and a solution exhibits ballistic propagation $d(t) = U_x t$ with

$$U_x = \gamma^{1/2} \times \left(\frac{\chi_0 I}{2} \right)^{1/2}$$

- $U_x \sim$ geometric mean of **“local growth”** and **“turbulent diffusion”**, faster than transport time scale.

Edge Turbulence Spreading to Unstable Core

- **Nonlinear Gyrokinetic Simulations of Ion Temperature Gradient Turbulence:**


$$\frac{R}{L_T} = 6.9 \text{ at core (Cyclone value)}$$

$$\frac{R}{L_T} = 13.8 \text{ at edge}$$


- Initial Growth at Edge followed by **Ballistic Front Propagation** into Core

- Saturation Level at Core $\sim 2\times$ Core (only) Result


$$\nabla \cdot \Gamma_I \sim \gamma_{local} I$$



Front Propagation Speed Increases with R/L_T




- From Simulation, U_x and I increase with $(\frac{R}{L_T})$
- Nonlinear Diffusion Model: $U_x \propto (\gamma I)^{1/2}$
by [Gurcan-Diamond-Hahm-Lin, *submitted to PoP '04*]



- Toroidal Linear Coupling dominant Regime: $U_x \sim \frac{\rho_i}{R} v_{Ti}$
by [Garbet-Laurent-Samain-Chinardet, NF '94]
- Four Wave Model: Complex Bursty Spreading
by [Zonca-White-Chen, PoP '04]

Summary

- Turbulence spreading has been widely observed in global gyrokinetic particle simulations: It can be responsible for deviation of transport scaling from GyroBohm.
- Fluctuation Intensity in the linearly stable region can be significant due to turbulence spreading.
- Sometimes **Spreading of Edge Turbulence** into Core can exceed local turbulence in connection region.
- It is likely to affect “the edge boundary conditions” used in core modeling, and predictions of pedestal extent.




Turbulence Spreading has been widely observed

- From Most Global Gyrokinetic/Gyrofluid Simulations:
 - X. Garbet *et al.*, NF '94 (Mode-coupling in Torus)
 - R. Sydora *et al.*, PPCF '96 (Torus with Zonal Flows)
 - Y. Kishimoto *et al.*, PoP '96 (Torus with Zonal Flows)
 - S. Parker *et al.*, PoP '96 (Torus without Zonal Flows)
 - W.W. Lee *et al.*, PoP '97 (Torus without Zonal Flows)
 - Y. Idomura *et al.*, PoP '00 (Sheared Slab with Zonal Flows)
 - **Z. Lin** *et al.*, **PRL '02** (Torus with Zonal Flows)
 - L. Villard *et al.*, IAEA '02 (Cylinder with Zonal Flows)
 - R. Waltz *et al.*, PoP '02 (Torus with Zonal Flows)
 - Y. Kishimoto *et al.*, H-mode '03 (Sheared Slab with ZF)
- **Neither** Zonal Flows nor Toroidal Coupling necessary for Turbulence Spreading.

Anomalous Transport where $\gamma_{lin} < 0$

Core of Reversed Shear Plasmas where profiles are nearly flat (JT-60U, TFTR, DIII-D,...)



→ Nonlinearly Unstable?
(Self-sustained Turbulence (B. Scott))

→ **Spreading from the Linearly Unstable Zone**

Distinction between “Core” and “Edge” blurred


- Researchers have frequently divided the tokamak into three zones — a central sawtooth zone, a middle ‘confinement zone’, and an edge zone...

Goldston-U.S.A. *Kyoto IAEA (1986)*

- the edge..., often used as a boundary condition for core transport modeling

V. Parail, *Plasma Phys. Control. Fusion*, **44**, A63 (2002)

- $\frac{\partial}{\partial x} \gamma(x) \sim \frac{\partial^2}{\partial x^2} P$: large at the top of pedestal



Long Term Behavior: Sub-Diffusion



- Self-similar Variable: $\ell(t)^2 \sim \chi_0 I^\beta t$
- $I(t)\ell(t) = I(0)\ell(0) \equiv \epsilon$, up to dissipation
- $\ell(t) \sim [\chi_0 \epsilon^\beta t]^{\frac{1}{2+\beta}}$
 - $\sim t^{1/3}$: Weak Turbulence
 - $\sim t^{2/5}$: Strong Turbulence
- Previous numerical mode coupling study:
 - X. Garbet *et al.*, NF 1994
 - Linear toroidal coupling usually dominates $\sim t^1$: convective
 - Without linear toroidal mode coupling $\sim t^{1/2}$: **diffusive**

Short Term Behavior: Ballistic Propagation

- $x_{front} = (x_0^3 + 6\epsilon\chi_0 t)^{1/3}$
- $U_x = \frac{d}{dt}x_{front}$
 - $\sim 2\epsilon\chi_0/x_0^2$: for small t (consequence of $\Delta \ll x_0$)
 - $\sim t^{-2/3}$: for large t (sub-diffusion)Note: $\epsilon \propto I$, turbulence intensity
- Scaling of U_x drastically different from V_{gr} of linear drift (ITG) wave
 - contrast our theory from others relying on linear dispersion
 - [eg., Garbet *et al.*, PoP '96; Zonca *et al.*, PoP '04]

Simple theory captures ρ^* dependence of spreading

$$I_0(x, t) = \frac{\epsilon}{(6\epsilon\chi_0 t + W^3)^{1/3}} \left(1 - \frac{(x - x_i)^2}{(6\epsilon\chi_0 t + W^3)^{2/3}} \right) \times H \left((6\epsilon\chi_0 t + W^3)^{1/3} - |x - x_i| \right)$$



Spreading of Self-sustained Turbulence

[Itoh, Itoh, Hahm, and Diamond, *submitted to J. Phys. Soc. Jpn.* '04]

$$\frac{\partial}{\partial t} I = \Gamma_{NL}(I, x)I + \chi_0 \frac{\partial}{\partial x} \left(I \frac{\partial}{\partial x} I \right)$$

Model self-sustained sub-critical turbulence [eg., B. Scott, *PRL* '90]:

$$\Gamma_{NL}(I, x) > 0 \text{ for } I_{crit} < I < \frac{\gamma_0}{\alpha},$$

$$\Gamma_{NL}(I, x) = 0 \text{ for } I < I_{crit}, I > \frac{\gamma_0}{\alpha}, \text{ at } |x| < L, \text{ and}$$

$\Gamma_{NL}(I, x) < 0$ at $|x| > L$ according to local linear and non-linear damping

Due to turbulence spreading, there exists a minimum size system (L) that can sustain the self-sustained turbulence