# Long Pulse Physics via International Stellarator Collaboration

G. A. Wurden & the US stellarator community\*

A presentation to the FESAC Strategic Planning Committee
Gaithersburg, Maryland
June 5, 2014

\*D.T. Anderson<sup>1</sup>, A.H. Boozer<sup>2</sup>, D. Curreli<sup>3</sup>, J.P. Freidberg<sup>4</sup>, D.A. Gates<sup>5</sup>, J.H. Harris<sup>6</sup>, S.R. Hudson<sup>5</sup>, M. Landreman<sup>7</sup>,S. Lazerson<sup>5</sup>, J.D. Lore<sup>6</sup>, G.H. Neilson<sup>5</sup>, D.A. Maurer<sup>8</sup>, N. Pablant<sup>5</sup>, A. Reiman<sup>5</sup>, D. Ruzic<sup>3</sup>, D.A. Spong<sup>6</sup>, J.N. Talmadge<sup>1</sup>, F.A. Volpe<sup>2</sup>, H. Weitzner<sup>9</sup>, G.A. Wurden<sup>10</sup>, M.C. Zarnstorff<sup>5</sup>

1 - University of Wisconsin

2 - Columbia University

3 – University of Illinois

4 – Massachusetts Institute of Technology

5 – Princeton Plasma Physics Laboratory

6 – Oak Ridge National Laboratory

7 – University of Maryland

8 – Auburn University

9 - New York University


10 - Los Alamos National Laboratory

# US Collaboration on overseas stellarators leverages substantial world investment


- Stellarator research addresses high priority Greenwald gaps
  - 3D plasma exhaust solutions compatible with high-performance core
  - Extending high-performance plasmas to long pulses.
- Stable commitment to long term collaborations with Germany and Japan will yield important physics that will help fill the gaps
  - Required for the design of stellarator FNSF and reactor concepts

<sup>\*</sup> See the Stellarator White Paper: Control of High-Performance Steady-State Plasmas: Status of Gaps and Stellarator Solutions

# LHD and W7-X, the world's two superconducting long-pulse stellarator experiments



National Institute for Fusion Science, Toki, Japan



Max Planck Institute for Plasma Physics, Greifswald, Germany

- W7-X begins operation in 2015, will have
  - SC magnets, high-power heating systems
  - Magnetic island divertor
  - 30 minute pulse at full power
- LHD has been operating since 1998
  - well equipped with heating, diagnostics
  - helical divertor
  - Achieved 1 hour long pulse at reduced power

## **Targeted research Opportunities on LHD**


- Support US developed X-ray Imaging Crystal Spectrometer (XICS) on LHD
  - Minimal investment to restart this collaboration \$0.3 million/year
- Argon Impurity transport studies
- Island physics studies
- Confinement studies at high density

US XICS diagnostic provides profile measurements of:

Ion Temperature Electron Temperature Poloidal Flow Velocity

Time resolution: 20ms, Spatial resolution: 2cm. Non-perturbative to the plasma (no NBI).

Time resolved ion temperature profiles From XICS on LHD




## W7-X: opportunities/challenges

- W7-X will become a world-class fusion facility.
  - ITER-like high-heat-flux and steady-state operation (before ITER)
  - Valuable supplement to tokamak program
- US contributions to W7-X (>\$10M) have made a real difference.
  - Added important capabilities (trim coils, scrapers, imaging . . .for plasma control and power exhaust management) beyond scope of IPP budget.
  - Strengthened position of W7-X, both inside and outside Germany
- US has earned partnership opportunity in W7-X ....
  - US contributions are deeply integrated into W7-X. Interest in/competition from other countries is growing.
  - Any US stellarator strategy will benefit from partnership in W7-X.

### 3D Magnetics on W7-X: Island Divertors

- Exhaust heat from the W7-X plasma is carried by hot particles flowing along 5/5 island chain
- Divertor targets (5-fold symmetry) are placed so as to intersect the islands
  - Field lines are mostly toroidal, so they intersect targets at a shallow angle.
- As bootstrap current changes, so does the magnetic geometry of the islands at the plasma edge
- US designed "scraper element" protects parts of each divertor
  - Science program studies the effect of this divertor structure



### **W7-X US Collaboration**


- The driving themes of the existing collaboration are: Edge plasma/divertor physics in 3D geometry, core transport in 3D, and plasma control in long pulse operation. This includes simulation/modeling, diagnostics, and planning for experiments.
- We have a seat on the W7-X council. Up to a dozen US scientists and engineers are presently working back/forth with IPP Greifswald, with increasing efforts coming as operations begin next year.
- Ongoing activities
  - Deploying an X-ray imaging crystal spectrometer (ion temp and rotation profiles)
  - Field line mapping with US supplied trim coils (modeling support)
  - IR imaging and analysis, optical access periscope for Scraper (heat-load mapping)
  - Design and build Test Divertor Unit Scraper Element (island divertor studies)

### **Exciting opportunities for expanded collaborations**

- Proposed areas for expansion of US partnership on W7-X
  - Formation of university/laboratory W7-X science partnership
  - TDU scraper element fabrication.
  - Detailed divertor/scraper imaging, fast data processing.
  - Energetic particle physics
  - Fluctuation measurements, turbulence simulation
  - Steady-state pellet injector preparation
- These research areas would further leverage W7-X into our overall long-pulse, edge/plasma, 3D divertor development strategy.


### Scraper element is a target for <u>future</u> expanded joint U.S. University and National Laboratory program in edge physics

Proposal: Fabricate scraper element



**TDU Scraper observation** and physics exploitation

All topics proposed for startup phase are relevant **Proposal: Support divertor** endoscope development



**National Laboratory goals** 

Scraper with endoscope offers multiple opportunities to exploit university strength in resonance with nat. labs

**University contributions** 

#### Magnetic Structure Optimization



#### Edge physics and exhaust



#### Plasma material interaction

- **Control of current profiles**
- Island control with Trim coils
- Impact on scraper heat flux of both
- **Joint VMEC efforts**
- **STELOPT & V3FIT**
- **Expertise in 3D equilibrium**

- Impact of scraper on exhaust
- Scraper heat and particle flux
- **SOL** width with scraper
- **EMC3-Eirene modeling**
- **Exhaust measurements**
- **Probe diagnostics**

- Scraper as high heat flux test surface
- Localized erosion/deposition
- Spectroscopy
- Atomic models for emission
- **Optical design expertise**


# International collaboration on stellarators is a high leverage opportunity that can fill Greenwald gaps

- Integral component of a coherent US stellarator program
  - Broad participation with Universities and laboratories
- US is already engaged in an exciting productive program
- Taking full advantage of new opportunities require added investment, \$10-12 million/year
  - e.g. 10% partnership on W7-X
  - Renewing targeted LHD collaboration
  - US participation is welcomed on both LHD and W7-X
- Urgency for action is high as W7-X starts operation in 2015

# **Backup Slides**



### We need to work within the W7-X Schedule



### **Operational phase 1 (OP-1)**

- inertially cooled divertor, only partial cooling of in-vessel components
- up to 10 s at 10 MW, 50 s at 1 MW heating power
- development of steady state scenarios

### **Operational phase 2 (OP-2)**

- high heat flux divertor, water cooling of in-vessel components
- steady state at 10 MW, 10 s at 18 MW heating power
- transition to high power steady state operation (30 min)



### One possible new effort: W7-X pellet fueling

### Stellarators heated with ECH exhibit hollow n(r)

- Serious risk in W7-X of edge density rising to ECH cut-off and limiting heating.
- Verified in 30 years of experiments on ATF, Heliotron-E, CHS, W7-AS, LHD
- Ware pinch, which transports particles to axis in tokamaks, vanishes for  $I_{pl} = 0$
- Solution: repetitive pellet fueling as demonstrated on ATF(1990).

#### Planning for W7-X pellet fueling

Staged capabilities:

5-10 pellets for TDU phase (20 s)

Continuous pellet injection for steady-state operation (>2017)

Synergy with ITER R&D (and on nearer term time-scale)

ORNL is developing continuous pellet system for ITER

Pinch effect much reduced in non-inductive tokamaks: turbulent pinch?


Strategy meeting of pellet experts from IPP, ORNL, and NIFS at IPP was in April 2013
 Multi-faceted international collaboration plan has emerged: more leverage.

# W7-X at the beginning of 2014



Max Planck Institute for Plasma Physics, Greifswald, Germany

# The LHD torus hall



National Institute for Fusion Science, Toki, Japan